



Thomas G. Gillespie, DVM, Diplomate, ABVP
Swine Health Management Specialty

210 East Wood Road
Rensselaer, IN 47978
(219) 866-6465
Fax (219) 866-3431
www.rssvet.com

2010 4-H Pig Vaccination Recommendations



We have some new combination vaccines that will make routine vaccination programs easier for most of you raising club pigs. The following is a comprehensive recommendation for all 4-H clients. If you have additional or specific needs, please ask and we will work out a plan for you.

Weaning (3-4 weeks of age)

CircoFlex (PCV2) + MycoFlex (Mycoplasma)

Mix these two vaccines together (1 cc of each) for just the number of pigs that you will be vaccinating and give a single 2 cc dose. Once mixed, the vaccine should be used within 4 hours. The remaining unmixed products are stable and usable until the expiration date on the bottle.

5-6 Weeks of age

Toxivac + Parasuis (Atrophic Rhinitis, Erysipelas, Haemophilus parasuis)
Give a 2 cc dose

8-10 Weeks of age

A.) *Toxivac + Parasuis (Atrophic Rhinitis, Erysipelas, Haemophilus parasuis)*
Give another 2 cc dose 2-3 weeks after the first dose

B.) *PneumoStar SIV (Influenza)*
Give a single 1 cc dose

All vaccines should be given in the neck with a clean needle. An 18x1" needle is sufficient for all of these pig ages.